



Stony Brook  
Medicine

---

---

---

---

---

---

---

---



Stony Brook Medicine

***Treating Pregnant Women's  
Tobacco Use and  
Dependence***

Todd R Griffin, MD, FACOG  
Chair  
Department of Obstetrics, Gynecology and  
Reproductive Medicine  
Stony Brook Medicine



Stony Brook University Hospital

---

---

---

---

---

---

---

---



Stony Brook Medicine

**FACULTY DISCLOSURES**

- I have no conflicts of Interest for this Presentation
- I have never smoked 😊

---

---

---

---

---

---

---

---

Stony Brook Medicine OBJECTIVES

- Describe two evidence-based interventions to help pregnant smoker's quit
- Assess an organization's guidelines to treat pregnant women's tobacco use with evidence-based treatment recommendations
- Discuss at least one gain and one challenge when treating pregnant women's tobacco users during the course of their pregnancy and postpartum

---

---

---


---

---

---

---

---




---

---

---

---

---

---

---

---

Stony Brook Medicine TREATING PREGNANT WOMEN'S TOBACCO USE AND DEPENDENCE

**Demographics:**

- Lower Socioeconomic Class
- Lower Educational Level
- Younger
- Unmarried
- Higher Parity
- Lower Level of Social Support
- Increased Incidence of Depression/Stress
- Internationally (WHO report) – Incidence of Woman smoking is increasing – especially in low-income high population countries

---

---

---

---

---

---

---

---


Stony Brook Medicine

TREATING PREGNANT WOMEN'S  
TOBACCO USE AND DEPENDENCE

***Treating Pregnant Women's Tobacco Use and Dependence***

Most important, potentially preventable cause of adverse pregnancy outcomes

- Placental Abruption
- Miscarriage
- Preterm Birth
- Low Birth Weight / Intrauterine Growth Restriction
- Stillbirth




---

---

---

---

---

---

---


---

Stony Brook Medicine

TREATING PREGNANT WOMEN'S  
TOBACCO USE AND DEPENDENCE

***Long-Term Adverse Effects:***

- Higher Rate of Attention Deficit Hyperactivity Disorder
- Increased Risk of Asthma
- Adverse Effects on the Immune System
- Possibly Increased Risk of Childhood Cancers




---

---

---

---

---

---

---

---


Stony Brook Medicine

TREATING PREGNANT WOMEN'S  
TOBACCO USE AND DEPENDENCE

***Association of Stillbirth and Tobacco Use:***

*Stillbirth Collaborative Research Network – NICHD (Mar 2006-Sept 2008)*

Matched Stillbirth and Live Births  
Women Self-Reported Smoking  
Checked Umbilical and Maternal Levels of Serum Cotinine  
Evaluated Risk of Stillbirth




---

---

---


---

---

---

---

---

 **Stony Brook Medicine** TREATING PREGNANT WOMEN'S  
TOBACCO USE AND DEPENDENCE

Self Reporting of Tobacco Usage and Still Birth	
	Stillbirth Risk (Odds Ratio)
Never Smoked	1.0
1-9 Cigs/Day	1.77
> 10 Cigs/day	2.17

Cotinine Concentrations (ng/ml)	
Negative	1.0
50 <sup>th</sup> percentile or less	2.04
> 50 <sup>th</sup> percentile	2.39

They noted a general dose-response effect – strengthening the biologic plausability of the association of tobacco use and stillbirth

Obstetrics & Gynecology Jan 2014 Vol 123; 118-124

---

---

---

---

---


---

---

---

---

---

 **Stony Brook Medicine** TREATING PREGNANT WOMEN'S  
TOBACCO USE AND DEPENDENCE

***Secondhand Smoke and Pregnancy:***  
Meta-Analysis – 19 studies

	Odds Ratio
Spontaneous Abortion	1.17 (0.88-1.54)
Stillbirth	1.23 (1.09-1.38)
Congenital Malformations	1.13 (1.01-1.26)

23% Increased risk of stillbirth and 13% increase risk of congenital malformation

Leonardi-Bee; Et al Pediatrics 2011; 127; 734-741

---

---

---

---

---


---

---

---

---

---

 **Stony Brook Medicine** TREATING PREGNANT WOMEN'S  
TOBACCO USE AND DEPENDENCE

***Environmental Tobacco Smoke on Perinatal Outcomes:***

- Retrospective Cohort Study
- 11,852 women
- Primary Outcome Measures:
  - Birthweight
  - Birth Length
  - Head Circumference
  - Stillbirth

Crane, Et al BJOG Mar 2011; 865-871

---

---

---

---

---

---

---

---

---


---

Stony Brook Medicine

TREATING PREGNANT WOMEN'S  
TOBACCO USE AND DEPENDENCE

**Results:**

Lower Mean Birth weight  
Smaller HC and Birth Length  
Stillbirth (OR 3.35)  
Preterm Birth (OR 1.87)  
Neonatal Sepsis (OR 2.96)



Crane, Et al BJOG Mar 2011; 865-871

---

---

---

---

---

---

---

---

Stony Brook Medicine

TREATING PREGNANT WOMEN'S  
TOBACCO USE AND DEPENDENCE




---

---

---

---

---

---

---

---

Stony Brook Medicine

TREATING PREGNANT WOMEN'S  
TOBACCO USE AND DEPENDENCE

**Smoking and Early Placental Pathology:**

- Examined placentas from 20 heavy smokers and 20 non-smokers undergoing TOP at 9-14 weeks:
- Villous Changes and Morphologic changes in the Trophoblast occur very early in pregnancy
- Tobacco cessation is best before pregnancy occurs or as early as possible.

Jauniaux, et al Obstetrics & Gynecology; May 1992; 79 (645-648)

---

---

---

---

---

---

---


---



Stony Brook Medicine

TREATING PREGNANT WOMEN'S  
TOBACCO USE AND DEPENDENCE

### Smoking Cessation and Pregnancy:




Pickett, Et al; Obstetrics &amp; Gynecology; Jan 2003, 101 (140-147)



Stony Brook Medicine

TREATING PREGNANT WOMEN'S  
TOBACCO USE AND DEPENDENCE

### Fluctuations in Treating Pregnant



Need repeated interventions to promote smoking cessation

Pickett, Et al; Obstetrics &amp; Gynecology; Jan 2003, 101 (140-147)



Stony Brook Medicine

TREATING PREGNANT WOMEN'S  
TOBACCO USE AND DEPENDENCE

### Smoking Cessation Techniques:

- Psychosocial Interventions
- Pharmacologic Interventions

2014 Goals

1. Quit Smoking
2. Quit Smoking
3. Quit Smoking




Stony Brook Medicine

TREATING PREGNANT WOMEN'S  
TOBACCO USE AND DEPENDENCE

**Psychosocial Interventions:**

- Counseling
- Health Education
- Feedback
- Incentive-Based Interventions
- Social Support
- Other (exercise)



Cochrane Review (2013)

---

---

---

---

---

---

---

---

Stony Brook Medicine

TREATING PREGNANT WOMEN'S  
TOBACCO USE AND DEPENDENCE

**Counseling:**

- Motivational Interviewing / Cognitive Behavior Therapy, Psychotherapy, Relaxation, Problem Solving Facilitation, etc.
- Face to Face, Telephone, Interactive Computer Programs, AV equipment

**Health Education:**

- Women provided information on risks of smoking. May or may not have received additional support (ex. Automated texts, etc.)

---

---

---

---

---

---

---

---

Stony Brook Medicine

TREATING PREGNANT WOMEN'S  
TOBACCO USE AND DEPENDENCE

**Feedback:**

- Mother receives information about fetal health – US monitoring, CO or Urinary Cotinine Levels

**Incentive Based Interventions:**

- Mother receives financial incentive – gift vouchers, etc.

**Social Support (Peers and/or Partner)**

---

---

---

---

---

---

---

---

Psychosocial Techniques for Smoking Cessation		
Technique	Positive Impact	No Change/Negative
Counseling	++	
Health Education	+/-	+/-
Feedback	+	
Incentive Based	+++ (BUT ONLY IF PROVIDED INTENSIVELY)	
Social Support	+ (peer support)	0 (partner Support)

---

---

---

---

---

---

---

---

Stony Brook Medicine

TREATING PREGNANT WOMEN'S TOBACCO USE AND DEPENDENCE

**Pharmacologic Interventions:**  
 Nicotine Replacement Therapy  
 Varenicline  
 Bupropion




---

---

---

---

---

---

---

---

Stony Brook Medicine

TREATING PREGNANT WOMEN'S TOBACCO USE AND DEPENDENCE

**Nicotine Replacement Therapy:**

- Cochrane Review (2012)
- "Insufficient evidence to support either the efficacy or safety of NRT used in pregnancy"
- Mixed fetal/infant outcomes
- Increased risk of C/S noted
- No better than placebo

---

---

---


---

---

---

---


---


**Stony Brook Medicine**

TREATING PREGNANT WOMEN'S  
TOBACCO USE AND DEPENDENCE

**Bupropion:**

- Limited studies in Pregnancy
- Most Animal and Human first trimester exposure studies do not demonstrate increase risk of fetal malformations
- 1 Retrospective Study – Increased risk of Left outflow tract Heart Defects if used around conception (OR – 2.6)



Cochrane Review (2012)

---

---

---


---

---

---

---

---


**Stony Brook Medicine**

TREATING PREGNANT WOMEN'S  
TOBACCO USE AND DEPENDENCE

**Meta-Analysis of Pharmacotherapy for Smoking Cessation in Pregnancy:**

- Reviewed 74 articles – Only 7 RCTs
- Only evaluated NRT
- Demonstrated an Increase in Abstinence Rate: 1.8 times higher
- Of the 7; only 3 used a placebo control – this subgroup showed no increased efficacy
- **Conclusion:** Further, larger RCTs needed

Myung, et al, BJOG 2012; 119 (1029-1039)

---

---

---


---

---

---

---

---


**Stony Brook Medicine**

TREATING PREGNANT WOMEN'S  
TOBACCO USE AND DEPENDENCE

**Practice Patterns of Ob/Gyns and Smoking Cessation:**

- ACOG Survey – 252 Respondents
- 88.1% screen at initial visit
- Only 59.9% f/u at subsequent visits
- 35% never communicate tobacco use during pregnancy to the Pediatrician
- Most Common Methods Used:
  - Recommend Abstinence
  - Recommend Reduction
  - Review Reasons for Quitting

Coleman, et al, J Addict Med 2014; 8 (14-24)

---

---

---

---

---

---

---


---

Stony Brook Medicine

TREATING PREGNANT WOMEN'S TOBACCO USE AND DEPENDENCE

**Barriers to Intervention:**

- Time
- Expected Patient Denial or Resistance to Treatment
- Lack of Referral Resources



Coleman, et al, J Addict Med 2014; 8 (14-24)

---

---

---

---

---

---


---

---

Stony Brook Medicine

TREATING PREGNANT WOMEN'S TOBACCO USE AND DEPENDENCE

**Resources Requested by Ob/Gyns to Improve Efforts**



Resource Requested	Mean self-reported preparedness (scale 0-4)
Screen pregnant women for tobacco use	3.8
Educate pregnant women about the effects of tobacco on their unborn baby	3.8
Educate postpartum women about the effects of tobacco on their infant	3.4
Screen postpartum women for tobacco use	3.4
Conduct brief intervention with pregnant women who smoke	3.2
Prescribe smoking cessation medications for nonpregnant women	2.9
Utilize resources to refer patients who need a tobacco cessation program	2.6
Prescribe smoking cessation medications for pregnant women	2.4

Coleman, Et al, J Addict Med 2014; 8 (14-24)

---

---

---

---

---

---

---

---

Stony Brook Medicine

TREATING PREGNANT WOMEN'S TOBACCO USE AND DEPENDENCE

**Conclusions:**

- Multiple Adverse Fetal and Neonatal Outcomes associated with Primary and Secondary Tobacco Exposure
- If Smoking beyond first trimester few quit
- Strategies using Incentives seem to be most effective
- Pharmacotherapy interventions need further research
- Ob/Gyns need further education, training and assistance to be successful

---

---

---


---

---

---

---


---



Stony Brook Medicine

TREATING PREGNANT WOMEN'S  
TOBACCO USE AND DEPENDENCE

Thank You!



---

---

---

---

---

---

---